



# NAD-dependent protein deacetylase sirtuin-2

Cat.No:DTD0385

## DESCRIPTION

<b>Name</b>	<b>NAD-dependent protein deacetylase sirtuin-2</b>
<b>Purification</b>	The soluble target protein NAD-dependent protein deacetylase sirtuin-2 was derived from secreted expression and was purified by affinity chromatography
<b>Purity</b>	>85% by SDS-PAGE
<b>Molecular weight</b>	44 kDa
<b>Expression System</b>	HEK293
<b>Species</b>	Human
<b>Physical Appearance</b>	Liquid
<b>Buffer</b>	1×PBS, 5% Glycerol, pH7.4
<b>Concentration</b>	0.500 mg/ml, BSA was used as standard and protein concentration was determined by Bradford
<b>Stability &amp; Storage</b>	For long term use, aliquot and store at -80°C (avoiding multiple freeze-thaws)

### SDS-PAGE & Western Blot



Lane 1: BSA (1.00 µg)  
 Lane 2: SIRT2(WT) protein (0.50µg)  
 M<sub>1</sub>: SDS-PAGE Marker  
 M<sub>2</sub>: Western Blot Marker  
 Using Anti-His antibody

**Thank you for choosing Detai Biological. We are dedicated to serving you!**  
**For Research Use Only**